


Journal of Poultry Sciences and Avian Diseases

Journal homepage: www.jpsad.com



Public Health Impact of Staphylococcal and Clostridial Infections of Poultry: A Comprehensive Review



Peyman Nakhaee¹ , Hafez Mohamed Hafez^{2*} 

¹ Department of Clinical Sciences, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

² Former Head of the Institute of Poultry Diseases, Free University Berlin, Germany

* Corresponding author email address: hafez@vetmed.fu-berlin.de

Article Info

Article type:

Review Article

How to cite this article:

Nakhaee, P., & Mohamed Hafez, H. (2025). Public Health Impact of Staphylococcal and Clostridial Infections of Poultry: A Comprehensive Review. *Journal of Poultry Sciences and Avian Diseases*, 3(1), 13-25. <http://dx.doi.org/10.61838/kman.jpsad.3.1.2>



© 2025 the authors. Published by SANA Institute for Avian Health and Diseases Research, Tehran, Iran. This is an open access article under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License.

ABSTRACT

Staphylococcus aureus causes staphylococcal food poisoning and several difficult-to-treat infections in humans. The occurrence and dissemination of methicillin-resistant *S. aureus* (MRSA) is crucial and well documented in various studies. MRSA is an increasing public health concern worldwide. In addition, *S. aureus* is resistant to commonly used antibiotics in poultry farms, which is a concern to public health because of the transmission of this bacteria after consuming poultry meat. Hence, these highlight the significance of antimicrobial and enterotoxigenic monitoring of *S. aureus* in food chains. *Clostridium perfringens* is a ubiquitous spore-forming anaerobic pathogen that causes broilers' clinical or subclinical necrotic enteritis. At the same time, in humans, it is the causal agent of foodborne diseases, frequently related to the consumption of chicken meat. Moreover, enterotoxin-producing *C. perfringens* has high zoonotic potential as well as serious public health concerns due to the emanation of foodborne intoxication. The high diversity and occurrence of *C. perfringens* and *S. aureus* strains indicate the need to carry out various plans to control *C. perfringens* and *S. aureus* associated with foodborne infections. Meanwhile, clinical importance is assisting in understanding the occurrence, source, reservoir, and evolution of antibacterial resistance of *C. perfringens* and *S. aureus* to establish the control of these pathogens.

Keywords: *C. perfringens*, Foodborne diseases, Poultry, Public health, *S. aureus*

Article history:

Received 17 July 2024

Revised 27 September 2024

Accepted 04 October 2024

Published online 01 January 2025

1 Introduction

Staphylococcus, a genus of Gram-positive bacteria, encompasses several species, with *Staphylococcus aureus* and *Staphylococcus epidermidis* being the most commonly implicated in poultry infections. These bacteria can cause a spectrum of diseases ranging from mild skin infections to more severe systemic diseases in the poultry population. Staphylococci are commonly found in the environment, including animals' and humans' skin and mucous membranes (1, 2). While many species of *Staphylococcus* exist, *Staphylococcus aureus* and *Staphylococcus epidermidis* are among the most frequently encountered in poultry environments (3-5). These bacteria can lead to various bird infections, including skin and soft tissue infections, respiratory tract infections, and systemic diseases (2, 3, 6, 7). *Staphylococcus* infections cause significant challenges to the poultry industry worldwide, affecting bird health and productivity (6, 8, 9). Diverse factors, including management practices, environmental conditions, and other predisposing diseases, influence the occurrence of *Staphylococcus* infections in poultry. In commercial poultry production systems, high stocking density, suboptimal or low ventilation, and incomplete sanitation and biosecurity can provide favorable conditions for the proliferation and transmission of *Staphylococcus* spp. among birds (7, 10, 11). Additionally, stressors such as transportation, handling, and high-density housing can compromise the immune function of poultry, increasing their susceptibility to bacterial infections (4, 12, 13).

Staphylococcal infections in poultry manifest in various forms, including dermatitis, cellulitis, bumblefoot, omphalitis (yolk sac infection), arthritis, and respiratory tract infections. These infections can lead to significant economic losses for poultry producers because of decreased growth rates, impaired feed conversion efficiency, increased mortality rates, and costs related to veterinary treatments and interventions (8, 9, 11, 14, 15). Moreover, *Staphylococcus aureus*, in particular, poses a notable public health concern due to its zoonotic potential (7, 9, 15-17). *Staphylococcus aureus* infection in human occurs through direct contact with infected birds or their contaminated products, leading to a range of infections, including skin and soft tissue infections, respiratory tract infections, and foodborne illnesses (3, 6, 15, 18). Furthermore, the emergence of antibiotic-resistant *Staphylococcus* strains in poultry populations raises additional concerns regarding the effectiveness of antibiotic therapies for both animal and human health (9, 15, 18-23).

Effective prevention and control strategies are paramount because of the multifaceted nature of *Staphylococcus* infections in poultry (24, 25). This necessitates implementing comprehensive biosecurity measures, including strict hygiene protocols, routine surveillance, appropriate antibiotic stewardship, and management practices to minimize stress and optimize flock health (6, 9, 15, 16, 25-28). By investigating these factors, poultry producers can mitigate the impact of *Staphylococcus* infections on bird welfare, productivity, and public health, thereby ensuring the sustainability and profitability of the poultry industry. *Staphylococcus aureus* (*S. aureus*) is a significant zoonotic pathogen that can cause disease in humans and animals (3, 29). It is widely distributed in nature and is present in air, water, and feed; it also exists on the surface of the human body, in the nasal cavity, on animal fur, and in the digestive tract, among other sites. *S. aureus* has been responsible for several infectious diseases, including tissue and skin infections, pneumonia, sepsis, mastitis, arthritis, and soft tissue infections (3, 5, 30-32).

Clostridium species are Gram-positive, rod-shaped bacteria that thrive in anaerobic conditions. A hallmark of these bacteria is their ability to form endospores, which are highly resistant to environmental stresses, including heat, desiccation, and disinfectants (33-35). This spore-forming capability allows *Clostridium* to persist in the environment, contributing to the difficulty in controlling infections within poultry systems. They are commonly found in soil, decaying organic matter, and the gastrointestinal tract of animals, where they play roles ranging from benign commensals to serious pathogens (34, 36). *Clostridium* infections represent a significant challenge in poultry production, affecting the health and productivity of commercial poultry flocks worldwide (37, 38). The causative agents are various species within the genus *Clostridium*, anaerobic, spore-forming bacteria known for their pathogenic potential and environmental resistance (34, 39). These infections can lead to severe diseases such as necrotic enteritis, gangrenous dermatitis, botulism, and ulcerative enteritis, each characterized by distinct clinical manifestations and substantial economic impacts (34, 40-42). The transmission of *Clostridium* infections occurs primarily through the ingestion of spores from contaminated feed, water, or litter. These spores can germinate and proliferate in the gut under favorable conditions, such as disrupted gut flora, dietary changes, or environmental stress. Horizontal transmission within flocks is common, especially in intensive production systems with high stocking densities (34, 43). *Clostridium*

infections such as Necrotic Enteritis (NE) have remarkable economic implications due to high mortality rates during outbreaks, which result in substantial financial losses, poor feed efficiency, lower growth rates, reduced egg production, increased expenses related to medications, veterinary care, and enhanced biosecurity measures and infected birds suffer from severe symptoms, raising ethical and welfare concerns. Clostridial diseases in poultry production have profound economic and welfare impacts, and high mortality rates during outbreaks can lead to substantial financial losses due to loss of production and the need for carcass disposal (34, 38, 44). Meanwhile, infected birds often exhibit poor feed efficiency, reduced growth rates, and lower egg production, affecting overall productivity and profitability (45-47). Other implications of clostridial infections in the poultry industry are increased expenses for medications, veterinary care, and enhanced biosecurity measures, contributing to the economic burden and significant suffering in affected birds, leading to ethical considerations and potential regulatory scrutiny (38, 48).

2 Prevalence and Risk Factors

Staphylococcus species are ubiquitous in poultry environments, with *S. aureus* being the most clinically significant. Other species, such as *S. epidermidis* and *S. hyicus*, are also encountered, although they are less pathogenic (16, 49, 50). Various factors, including geographical region, type of poultry, age, and production stage, can influence the prevalence of *Staphylococcus* species in poultry. Various studies demonstrated a widespread prevalence of *S. aureus* and other *Staphylococcus* species in poultry farms across various regions, often exceeding 50% in some areas. A global review reported that *S. aureus* prevalence ranges from 20% to 80% in commercial poultry operations. (2, 6, 18, 51-53). Due to their intensive farming conditions, broilers often exhibit higher prevalence rates of *Staphylococcus* infections. In another study, several studies reported 30% to 80% prevalence rates depending on farm management and biosecurity practices (5, 9, 16, 30, 54). Researchers reported that the carriage rate of *S. aureus* among broilers on all farms was 84.8%, whereas it was 84% among farm workers. The differences in the incidence rates of *S. aureus* in broilers and broilers farm workers in all farms were statistically non-significant (55). In poultry, the most common type of infection is tenosynovitis and arthritis (7, 56). *Staphylococcus* infections occur mainly during the following four periods of a breeder's life. Omphalitis and femoral head necrosis often

relate to egg or hatchery contamination and minor surgeries during the first two weeks of life. Other lesions related to the *staphylococcus* infection are infected hock and stifle joints secondary to coccidiosis or harsh vaccine reactions. The milder forms of gangrenous dermatitis are generally caused by *S. aureus* (7, 57, 58). The organism must enter the circulatory system to cause disease; thus, any injury or lesion increases the possibility of disease. The most evident route of infection is through a break or injury in the skin, respiratory tract, and gut (7, 9, 58, 59). *S. aureus* has been able to adapt rapidly to several antibiotics, leading to the emergence of methicillin-resistant *Staphylococcus aureus* (MRSA). MRSA can resist various types of antibiotics, like β -lactams and others. After a decade, MRSA had been found in many countries that had been considered as an endemic in the mid-1970s (1, 6, 18, 60-65). The detection frequency of 25.9% on turkey farms relates to the results of the National Zoonosis Monitoring 2010, which found that 19.6% of farms were positive (7, 66, 67). In a regional study in the southwest of Germany, researchers found that 90% of the turkey farms were positive for MRSA (68-70). Limited data on the prevalence and epidemiology of MRSA in broiler flocks have been available so far. Out of 384 dust and fecal samples originating from broiler fattening farms, only 0.7% were suspected to be MRSA positive, and they discovered a significant proportion of diseased flocks (55, 71, 72). Other studies in different countries also had varying results, with 35% of flocks positive at slaughter in Denmark and 4 of 50 flocks positive in the Netherlands (61, 62, 65). In another study, the researchers found that 2 of 14 Belgian farms were positive for MRSA by sampling five broiler chickens at each farm (65, 73). Meanwhile, MRSA has been detected in chickens and turkeys from farms and slaughterhouses (12, 60-62, 74-76). LA-MRSA is a distinct spread of specific MRSA clones (ST-398) that colonizes various food animal species (including livestock and poultry) and may be associated with human infection. Risk groups and modes of transmission of livestock-associated MRSA (LA-MRSA) include persons with direct contact with MRSA colonized or infected livestock, such as veterinarians, meat vendors, farmers, workers at slaughterhouses, contact with many different animal products (62, 64, 77). LAMRSA isolates have been detected in meat in different geographic areas, raising concerns about the possibility of transmission of MRSA from the farm to the fork (12, 61, 62, 76). However, human MRSA strains (human-associated and community-associated MRSA) have also been discovered in poultry meat, indicating inappropriate hygienic and sanitary conditions

during slaughter or meat processing from poultry meat (6, 62, 78).

C. perfringens could cause reduced production performance in chickens. The intestinal function is to attain optimal feed conversion ratio (FCR) in which several factors reduce this index, such as the necrotic gut lesion and abnormal clostridium dominance in gut microflora together with clostridium toxins, all this reduced productivity (45, 79-81). Immunosuppression predisposes poultry, especially broiler chickens, to NE; immunosuppression likely changes the intestinal environment and commensal flora population (82-84). The recent shift to no antibiotics ever production has elevated the prevalence of Clostridium-related disease. The removal of antibiotic growth promotors (AGPs) has increased the mortality in no antibiotic ever broiler production by 25 to 50%, compared to conventional production (38, 85). With a high incidence of clostridial disease and mortality rates, it is vital to comprehend the prevalence and virulence specifications of *C. perfringens* within broiler flocks. Most diseases caused by *C. perfringens* are mediated by one or more toxins (86). *C. perfringens* is classified into seven toxigenic types, A to G, based on the production of six major toxins: alpha (α), beta (β), epsilon (ϵ), iota (ι), enterotoxin (CPE), and NetB (86-88). All toxigenic types of *C. perfringens* produce the α toxin encoded by the *cpa* gene. Additionally, type B produces β and ϵ toxin; type C produces β toxin; type D produces ϵ toxin; type E produces ι toxin; type F produces CPE, and type G produces NetB (38, 79, 80, 82, 84, 86-88). One or more *C. perfringens* toxins mediate these diseases. Gastrointestinal infections in humans and animals have been indicated to be related to *C. perfringens* type C. In contrast, other toxins have been demonstrated to cause disease in humans or animals, but not both. Type A is the most frequent of seven *C. perfringens* toxinotypes. However, type F is the one that causes food-related poisoning in humans (38, 80, 89). NE occurs in broiler chickens between the ages of 2 to 6 weeks and in layers of 12–24 weeks due to the reduced titer of maternal antibodies to the clostridial infections with clinical or subclinical manifestations. Outbreak of NE is frequently detected in broiler flocks kept on deep litter, which may be a source of infection (34, 41, 43, 82, 90-93). Clinical NE in poultry is identified by sudden increased mortality without premonitory signs, whereas a subclinical form is frequently related to poor weight gain and feed conversion ratio (47, 88, 91). The development of NE is not exclusively created by infection with *C. perfringens*; predisposing factors, including diets high in non-starch polysaccharide grains and fish meal protein, as well as protozoal infection, have been

demonstrated to have an impact on the incidence of NE (93-95). Over the years, antimicrobial agents have been used as growth promotors known as antibiotics growth promotors (AGP) and to control NE in poultry (82, 90, 92, 94-97). However, the prohibition of the usage of AGPs by the European Union in 2006, strict regulations, and voluntary withdrawal of AGPs to struggle with antibiotic-resistant bacteria strains and high consumer demand for antibiotic-free meat had led to the recurrence of NE in poultry (82, 92-94).

3 Antimicrobial Resistance

Methicillin, a new antibiotic with resistance to β -lactamase, was developed two decades later, in 1960, and in less than two years, the first methicillin-resistant *S. aureus* (MRSA) appeared. The term LA-MRSA is distinguished by the spread of certain MRSA clones (ST-398) that colonize various food animal species (including livestock and poultry) and may be associated with human infection (62-64). Risk groups and modes of transmission of livestock-associated MRSA (LA-MRSA) include persons with direct contact with MRSA colonized or infected livestock, such as veterinarians, meat vendors, farmers, workers at slaughterhouses, contact with many diverse animal products and transporters of livestock (60-62, 64, 98). *S. aureus* can adapt rapidly even during selective antimicrobial pressure, which caused the emergence of methicillin-resistant *S. aureus* (MRSA). Resistance to methicillin and other β -lactam antibiotics is due to the *mecA* gene or its *mecC* homolog, located in a mobile genetic element called the staphylococcal chromosomal cassette (65, 76, 99). MRSA has become a threat to public health because, globally, it causes infections associated with hospitals, general communities, and LA-MRSA (100, 101). Various antimicrobial agents such as β -lactamases, macrolides, aminoglycosides, and tetracyclines are widely used in poultry flocks for the prevention and treatment of staphylococci and other infections, which result in the development and emergence of drug-resistant strains of bacterial microorganisms (7, 22, 64, 102, 103). Another significant challenge facing human health is antibiotic resistance. The occurrence of antibiotic-resistant foodborne pathogens is also increasing because of their excessive use in human and animal treatments (14, 15, 18, 23, 104). Moreover, the determinant factors of antibiotic resistance can be transferred to other pathogenic bacterial agents, which can compromise the treatment of serious bacterial infections and, thus, create a great threat to public health (14, 23, 50, 99, 103, 104).

The prevalence of MRSA in animal samples from the prescreened farms averaged between 61.7% and 80% in turkeys and 50% and 54.2% in boilers, respectively. In a study, researchers identified 0% and 28% MRSA-positive animals on three broiler farms (51, 105). Other studies revealed a prevalence of only 6.9% in chickens at abattoirs and 4.4% after investigating pooled throat swabs from multiple broiler farms (6, 64, 106). In analyzing dust and fecal samples from broiler farms in Germany, only 0.7% were suspected to be MRSA-positive (71). Studies in other European countries also had different results, with 35% of flocks positive at slaughter and 8% during rearing (18, 63, 73, 99, 100, 107). In another study, the researchers found that 2 of 14 Belgian farms were positive for MRSA by sampling five broilers at each farm (108). Several studies demonstrated more resistance of *S. aureus* isolates to ciprofloxacin, 92.9% and 50% resistance of tetracycline and ampicillin, respectively. Meanwhile, other researchers reported 100% resistance in *S. aureus* isolates from poultry meat against tetracycline and 61.5% against methicillin in Nigeria (10, 109). They stated 46.2% and 15.4% resistance against chloramphenicol and ciprofloxacin, whereas 38.5% against gentamicin and sulfamethoxazole/trimethoprim. Multidrug-resistant *S. aureus* has been reported several times (78, 110, 111). In addition, several studies reported 59.2% tetracycline to *S. aureus* isolates from ready-to-eat food (112-114). Extensive antibiotic use is thought to be the major cause of drug resistance in foodborne pathogens. The high rate of antimicrobial susceptibility may be due to the low use of these antimicrobials in layer breeders compared with broiler breeders or broilers. Although *S. aureus* is implicated in human food poisoning, most poultry strains do not produce the enterotoxins that cause human foodborne disease (18, 107, 108, 110, 111, 113, 115).

With extensive antimicrobial agents, especially antibiotics, *S. aureus* resistance has indicated an increasing trend recently, which brings great challenges for the clinical treatment of infectious diseases (18, 50, 100, 104, 116). MRSA is a remarkable concern that causes severe morbidity and mortality worldwide (14, 18, 30, 31, 50, 51, 60, 61, 63, 64, 73, 99, 100, 104, 115, 116). More and more molecular epidemiological studies have demonstrated that there are diverse sources of MRSA strains, including hospital-associated MRSA (HA-MRSA), community-associated MRSA (CA-MRSA), and LA-MRSA, indicating that continuous surveillance and screening are essential to discovering modifications in the epidemiology of MRSA infection in humans and animals (18, 73, 99, 107, 117). *S.*

aureus is an opportunistic pathogen capable of causing severe disease to farm workers and domesticated animals (14, 55, 118). *S. aureus* is prone to obtaining resistance to many kinds of antibiotics. In the early 1940s, the concept of resistance of *S. aureus* strains to antibiotic therapy was raised, and the prevalence of antibiotic resistance has significantly increased in recent decades because of the excessive use of antibiotics and the prescription used for therapeutics of diseases (14, 55, 118). Since 1959, penicillin-resistant *S. aureus* infections have been successfully treated with methicillin. However, in 1961, several reports from the United Kingdom demonstrated that *S. aureus* strains had been resistant to methicillin. This was the first discovery of the methicillin-resistant *S. aureus* strains (18, 106). *S. aureus* isolated from poultry has become a serious zoonotic risk factor on a global scale for the farm workers groups who handle or live in close proximity to chickens (6, 13, 55, 60, 62). An increasing series of evidence proved that livestock workers are a high-risk population for LA-MRSA carriage because of regular close contact with animals, especially in farm-intensive areas (60, 99). The primary result was LA-MRSA colonization among study participants who had close contact with livestock, identified by LA-MRSA strain isolated from nasal, oropharyngeal, or axillary samples (18, 60, 61, 64, 73, 119, 120). The risk of developing LA-MRSA colonization among livestock workers and veterinarians is also influenced by other risk factors, including livestock density and the type of farm or stage of animal production, history of antimicrobial consumption, hospital admission, working on farms with MRSA-positive animals, flock size and sanitary conditions (101, 119-121). Currently, MRSA strains have been identified using the biochemical test to detect the phenotypical characterization of MRSA, which showed resistance to all the antibiotics that have been produced.

A very significant challenge is the continuous antibiotic resistance of *C. perfringens* strains. Not only is antimicrobial use associated with high antibiotic resistance among bacterial microorganisms, but selection due to utilization of antibiotics may also contribute to a positive or negative relationship with virulence determinants (39, 122, 123). Resistance and virulence may not frequently be independent properties, and their relationship may play a vital role in the pathogenesis of *C. perfringens* infection (38, 80, 122, 123). AGPs were the major intervention strategy against Clostridium infections in commercial broilers (38, 41, 44, 80, 87, 124, 125). This therapeutic protocol supports birds by direct bactericidal and bacteriostatic effects, modifying the gut microbiota, decreasing GIT inflammation, and improving the overall

physical health of the GIT (45, 47, 80, 86, 112, 122, 126). However, due to concerns about occurrence of antimicrobial resistance, poultry farmers and veterinarians have limited their antibiotic usage. In 2011, the annual Agricultural Resource Management Survey inferred that 48% of grow-out operations raised poultry, especially broilers, without antibiotics usage and only provided antibiotics when birds suffered from bacterial infections (123, 126-129). Recently, it was estimated that more than 50% of the industry raises broilers without antibiotics. This type of broiler production is referred to as no antibiotics ever (NAE). Broilers reared within NAE facilities cannot receive antimicrobials in feed, water, supplementation, or injection at any part of the bird's lifetime. (130-132). The potential increase in the antimicrobial resistance of *C. perfringens* has occurred recently, with several reports declaring that most *C. perfringens* were multidrug-resistant (MDR) isolates (39, 133, 134). The resistance to tetracycline through TetA(P) protein, which regulates tetracycline active efflux, was common. In addition, over 50% of *C. perfringens* isolates were resistant to lincomycin. However, 25% of this resistance was related to the expression of the *lnu* gene (39, 134, 135). It should be noted that higher minimum inhibitory concentration (MIC) values of amoxicillin and ciprofloxacin were recorded due to the presence of the β -lactamase (*bla*) and quinolone (*qnr*) resistance genes, respectively. In the same context, the macrolide-resistant *C. perfringens* may act as reservoirs for the *erm* gene, which assists in its conjugal transfer (39, 122, 123, 135-138).

Previous studies have shown that tetracycline resistance is the most common antimicrobial resistance phenotype observed in *C. perfringens* and that it is related to the use of antibiotics for the treatment and as growth promoters in food animals. As a result, antimicrobial resistance of *C. perfringens* to tetracycline, lincomycin, and erythromycin has increased significantly over the past three decades (20, 28, 39, 45, 111, 122, 123, 139, 140). *C. perfringens* induces a toxicoinfectious foodborne sickness, in which infection by viable bacterial cells and their toxins plays a significant role in creating gastroenteritis in the host (34, 141). *C. perfringens* infections cause gas gangrene, necrotizing enteritis, and food poisoning in humans and animals. In order to minimize and decrease the economic losses induced by these infections, many antimicrobials, such as ampicillin, tetracycline, chloramphenicol, metronidazole, and imipenem, have been used preventively in the livestock industry (19, 22, 39, 80, 89, 103, 134, 135). Despite increasing awareness of the importance of combating antimicrobial resistance to improve public health, resistance of *C. perfringens* from various

sources to different antibiotics has yet to be actively investigated. The long-term and frequent use of antibiotics in the livestock and poultry industry has increased antimicrobial resistance in *C. perfringens*, compromised the efficacy of antibiotics, and generated great difficulties in clinical treatment (20, 39, 122, 139, 140). Probiotics are extensively used microorganisms to deal with particular diseases such as avian subclinical NE. The efficacy of probiotics belonging to the genera *Bacillus*, *Lactobacillus*, *Enterococcus*, *Bifidobacteria*, and *Saccharomyces* has been assessed both in vivo and in vitro (80, 92, 96). One of these studies reported a meta-analysis that included independent trials performed in different countries simultaneously, demonstrating in large-scale assessments that the supplementation of probiotics like *B. subtilis* remarkably elevates productive parameters and decreases the histological damage caused by *C. perfringens* (44, 92, 124, 125, 132, 134, 136, 142-144). The composition of the microbiome associated with broilers has been associated with improved production efficiency, representing that the use of probiotics demonstrated a viable alternative to avoid antibiotics in diets. It has been suggested that probiotics may beneficially affect the structure of the host gut microbiota, consequently improving the growth and survival of farm organisms (44, 92, 125, 128, 132, 136, 142-144). In addition to probiotics, other strategies, including prebiotics, synbiotics, organic acids, phytochemical additives, and dietary modifications and enzymes, can be used for the prevention and control of *C. perfringens* in the poultry industry and, more importantly, avoid the occurrence of antibiotic resistance.

4 Public Health Impacts

Foodborne diseases (FBDs) and poisoning are the widespread and great public health challenges of the modern world. Both developed and developing countries are widely affected by foodborne infections. FBDs impact human health and well-being and have economic effects on individuals and countries. Food poisoning outbreaks primarily comprise meat and meat products, but other food items, such as milk, may also be contaminated (145-147). FBDs are pathological cases created by the ingestion of food containing biological, chemical, or physical hazards. Multiple pathogenic bacteria can cause FBD, among which *Salmonella* spp., *Staphylococcus aureus* (*S. aureus*), and *C. perfringens* are the most common pathogenic bacteria in poultry-source foods. (8, 139, 148).

Staphylococcal food poisoning (SFP) is in relationship with emetic activity, sepsis-related infections, pneumonia, and

toxic shock syndrome (TSS). *S. aureus* has been frequently isolated from many foodstuffs, such as dairy products and meat. Therefore, it has been accounted as the third largest cause of food-associated disorders worldwide (128, 139, 145). In addition, it is one of the major public concerns since the treatment protocol for infections is a challenge when facing resistance, contributing to the evolution of MRSA. MRSA from poultry meat poses a public health risk that can be transmitted to humans by handling or consuming contaminated poultry meat (127, 139, 140). MRSA in poultry and poultry products is a concern to the poultry industry because of the risk to human health (13, 50, 78, 115, 147, 149, 150). The most important concern is the production of staphylococcal enterotoxins by MRSA strains, which can induce staphylococcal foodborne illness (18, 51, 62, 73, 100, 107). Raw meat handling, cross-contamination, and undercooked meat consumption may lead to MRSA infections. The epidemiology of MRSA in poultry has gained significance in recent years because of the growing demand for high-quality food and improved food safety (151, 152). This demand has led to increased poultry production, processing, and distribution of products, facilitating MRSA creation and spreading all over the food production chain.

The spread of MRSA in livestock is a serious public health threat. MRSA occurs among livestock animals like pigs, cattle, poultry, and companion animals. Interestingly, LA-MRSA also emerged among humans, demonstrating a zoonotic transmission from animals to humans (61, 62, 147, 153). Working and/or living on an MRSA-positive broiler farm was identified as a risk factor for acquiring MRSA. According to previous studies conducted in European countries, approximately 35% of slaughterhouses were MRSA-positive. On the other hand, the researchers recorded a 28% prevalence of MRSA from poultry farms in Malaysia (51, 62, 154). Furthermore, they isolated MRSA from turkey flocks in the Netherlands at 62-80% (6, 73). In Germany, transmission of MRSA from livestock to humans occurred mainly from occupational animal contact (61, 62, 151). Among 466 persons tested for MRSA in Dutch poultry slaughterhouses, 26 individuals were positive, which specifies a higher exposure risk to MRSA compared to the non-occupational Dutch people (155). Furthermore, MRSA has been discovered in various meat products, including turkey, raw chicken, veal, pork, mutton or lamb, beef, and rabbit (3, 18, 154, 156-158). Many authors have referred to detecting MRSA in chickens, cattle, pigs, and dogs. Along with creating cultural awareness of humans worldwide, consumers tend to eat low-fat with high minerals, vitamin contents, good quality

protein, quickly prepared, and low expensive chicken meat compared to the other types of meat. However, at the same time, human exposure to food poisoning was increased by consuming contaminated chicken meats with MRSA (3, 9, 14, 18, 64, 73, 158).

Evaluating *S. aureus* prevalence in poultry flocks is important for future risk prediction in poultry production and related occupational risks. Hence, understanding the possible outbreaks in antibiotic resistance through a sustainable surveillance program of the antimicrobial resistance profile of *S. aureus* simultaneously with the medical and veterinary conditions is necessary for the appropriate improvement of *S. aureus* control (18, 149, 154). Regular genotypic analysis of *S. aureus* from human and chicken origin is needed to discover the relationship between them due to the potential for transmitting MRSA to humans by consuming poultry or any by-product containing such *S. aureus* strains. Despite decreased antimicrobial usage in European animal production in recent years, the prevalence of MRSA in farm animals has not reduced (61, 108). It is, therefore, likely that MRSA remains in other reservoirs, such as in humans (i.e., farm workers and veterinarians). Strategies involving reducing the reliance on antibiotic usage and the frequent change of antibiotic classes should be considered for reducing MRSA levels in highly contaminated farms. Control and treatment strategies such as strict preventive biosecurity measures, selective probiotic feeding, and MRSA eradication programs should be used to prevent MRSA outbreaks on a farm, reduce MRSA carriage and eradicate MRSA from contaminated farms, respectively (9, 61, 64, 108, 109, 132, 152, 159).

Approximately 13% of gastrointestinal foodborne outbreaks have been related to *C. perfringens* infections (80, 89, 134). *C. perfringens* foodborne infections are frequently associated with meat and poultry products. The meat products can be contaminated with this pathogen during slaughtering through the contaminated surface or the contact of carcasses with feces (80, 141, 160, 161). Standard food service practices should be pursued to prevent this pathogenic microorganism (82, 84, 134). Chicken meat is the most consumed animal protein, and a sufficient supply for consumers requires mass production strategies, exacerbating the challenge of infections by FBPs such as *C. perfringens*. Because of this, there is a need to find economical, environmentally friendly, and efficient alternatives in the modulation of the intestinal microbiota, which contribute to the efficient production of broiler chicken to meet current and future demand (41, 80, 162). Also, it is recommended to cook food until the internal temperature reaches 70°C. Notably, gastrointestinal infection

with *C. perfringens* in animals and humans occurs due to the production of potent exotoxins (89, 134, 136, 163). As one of the most common foodborne zoonotic pathogens, *C. perfringens* is known to induce a variety of diseases, from food poisoning and slight diarrhea to life-threatening enterotoxaemia and gas gangrene in humans and animals (80, 162-164). Due to the increase in the emerging threat of foodborne-associated *C. perfringens* infections, many investigations demonstrated the prevalence of *C. perfringens* in food chains. They highlighted the evolution hazards of *C. perfringens* and the widespread MDR/toxigenic phenotypes. Several studies have revealed the high prevalence of MDR *C. perfringens*, which is a remarkable threat to the efficient treatment of foodborne illness by restricting the therapeutic options (34, 39, 134, 165, 166). The results of related studies indicated frustrating susceptibility patterns, and the antibiotics used in poultry feed as growth promoters were the main reasons for the evolution of *C. perfringens* resistance patterns as the bacteria became adapted due to the frequent use of antibiotics. Hence, there is an essential need for strict guidelines explaining the use of antibiotics and the safe production of food products of poultry origin (39, 89, 122, 134, 136, 138, 165, 166).

Acknowledgements

None.

Conflict of Interest

The authors declared no conflicts of interest.

Author Contributions

All the authors contributed to all parts of this research.

Data Availability Statement

Data are available from the first author upon reasonable request.

Ethical Considerations

There is no ethical consideration.

Funding

This research has no grant from any public, commercial, or not-for-profit funding agency.

References

1. Fayisa WO, Tuli NF. Review on Staphylococcus aureus. Int J Nurs Care Res. 2023;1:1-8.
2. Frana TS, Hau SJ. Staphylococcosis. Diseases of swine. 2019;926-33. [DOI]
3. Haag AF, Fitzgerald JR, Penadés JR. Staphylococcus aureus in Animals. Microbiology spectrum. 2019;7(3):10.1128/microbiolspec.gpp3-0060-2019. [PMID: 31124433] [PMCID: PMC11257167] [DOI]
4. Li Q, Li Y, Tang Y, Meng C, Ingmer H, Jiao X. Prevalence and characterization of Staphylococcus aureus and Staphylococcus argenteus in chicken from retail markets in China. Food Control. 2019;96:158-64. [DOI]
5. Sanz S, Olarte C, Hidalgo-Sanz R, Ruiz-Ripa L, Fernández-Fernández R, García-Vela S, et al. Airborne dissemination of bacteria (enterococci, staphylococci and enterobacteriaceae) in a modern broiler farm and its environment. Animals. 2021;11(6):1783. [PMID: 34203681] [PMCID: PMC8232102] [DOI]
6. El-Ghany WAA. Staphylococcus aureus in poultry, with special emphasis on methicillin-resistant strain infection: a comprehensive review from one health perspective. 2021. [DOI]
7. Szafraniec GM, Szeleszczuk P, Dolka B. Review on skeletal disorders caused by Staphylococcus spp. in poultry. Veterinary Quarterly. 2022;42(1):21-40. [PMID: 35076352] [PMCID: PMC8843168] [DOI]
8. Hafez HM, Attia YA. Challenges to the poultry industry: Current perspectives and strategic future after the COVID-19 outbreak. Frontiers in veterinary science. 2020;7:516. [PMID: 33005639] [PMCID: PMC7479178] [DOI]
9. Szafraniec GM, Szeleszczuk P, Dolka B. A review of current knowledge on Staphylococcus agnetis in poultry. Animals. 2020;10(8):1421. [PMID: 32823920] [PMCID: PMC7460464] [DOI]
10. Akbar A, Anal AK. Prevalence and antibiogram study of Salmonella and Staphylococcus aureus in poultry meat. Asian Pacific journal of tropical biomedicine. 2013;3(2):163-8. [PMID: 23593598] [DOI]
11. Kierończyk B, Rawski M, Józefiak D, Świątkiewicz S. Infectious and non-infectious factors associated with leg disorders in poultry—a review. Annals of animal science. 2017;17(3):645-69. [DOI]
12. Fluit A. Livestock-associated Staphylococcus aureus. Clinical Microbiology and Infection. 2012;18(8):735-44. [PMID: 22512702] [DOI]
13. Govender V, Madoroba E, Magwedere K, Fosgate G, Kuonza L. Prevalence and risk factors contributing to antibiotic-resistant Staphylococcus aureus isolates from poultry meat products in South Africa, 2015-2016. Journal of the South African Veterinary Association. 2019;90(1):1-8. [PMID: 31478731] [DOI]
14. Pal M, Gutama KP, Koliopoulos T. Staphylococcus aureus, an important pathogen of public health and economic importance: A comprehensive review. Journal of Emerging Environmental Technologies and Health Protection. 2021;4(2):17-32.
15. Pal M, Kerorsa GB, Marami LM, Kandi V. Epidemiology, pathogenicity, animal infections, antibiotic resistance, public health significance, and economic impact of staphylococcus aureus: a comprehensive review. American Journal of Public Health Research. 2020;8(1):14-21.
16. Hossain J, Islam S, Sobur A, Zaman SB, Nahar A, Rahman M, et al. Exploring poultry farm environment for antibiotic resistant Escherichia coli, Salmonella spp., and Staphylococcus spp. having public health significance. 2020. [DOI]
17. Wibisono FM, Wibisono FJ, Effendi MH, Plumeriastuti H, Hidayatullah AR, Hartadi EB, et al. A review of salmonellosis on poultry farms: Public health importance. Systematic Reviews in Pharmacy. 2020;11(9):481-6.

18. Algammal AM, Hetta HF, Elkelish A, Alkhalifah DHH, Hozzein WN, Batiha GE-S, et al. Methicillin-Resistant *Staphylococcus aureus* (MRSA): one health perspective approach to the bacterium epidemiology, virulence factors, antibiotic-resistance, and zoonotic impact. Infection and drug resistance. 2020;3:255-65. [PMID: 33061472] [PMCID: PMC7519829] [DOI]
19. Aslam B, Khurshid M, Arshad MI, Muzammil S, Rasool M, Yasmeen N, et al. Antibiotic resistance: one health one world outlook. *Frontiers in cellular and infection microbiology*. 2021;11:771510. [PMID: 34900756] [PMCID: PMC8656695] [DOI]
20. de Mesquita Souza Saraiva M, Lim K, do Monte DFM, Givisiez PEN, Alves LBR, de Freitas Neto OC, et al. Antimicrobial resistance in the globalized food chain: A One Health perspective applied to the poultry industry. *Brazilian Journal of Microbiology*. 2022;1-22. [PMID: 34775576] [PMCID: PMC8590523] [DOI]
21. Heaton CJ, Gerbig GR, Sensus LD, Patel V, Smith TC. *Staphylococcus aureus* epidemiology in wildlife: A systematic review. *Antibiotics*. 2020;9(2):89. [PMID: 32085586] [PMCID: PMC7168057] [DOI]
22. Mak PH, Rehman MA, Kiarie EG, Topp E, Diarra MS. Production systems and important antimicrobial resistant-pathogenic bacteria in poultry: a review. *Journal of Animal Science and Biotechnology*. 2022;13(1):148. [PMID: 36514172] [PMCID: PMC9749270] [DOI]
23. Rao S, Linke L, Magnuson R, Jauch L, Hyatt DR. Antimicrobial resistance and genetic diversity of *Staphylococcus aureus* collected from livestock, poultry and humans. *One Health*. 2022;15:100407. [PMID: 36277090] [PMCID: PMC9582408] [DOI]
24. Bisola M-AI, Olatunji G, Kokori E, Mustapha AA, Scott GY, Ogieuh IJ, et al. Emerging challenges in innate immunity: *Staphylococcus aureus* and healthcare-associated infection. *Journal of Medicine, Surgery, and Public Health*. 2024;100103. [DOI]
25. Endale H, Mathewos M, Abdeta D. Potential Causes of Spread of Antimicrobial Resistance and Preventive Measures in One Health Perspective-A Review. *Infection and Drug Resistance*. 2023;7:515-45. [PMID: 38089962] [PMCID: PMC10715026] [DOI]
26. Butucel E, Balta I, McCleery D, Morariu F, Pet I, Popescu CA, et al. Farm biosecurity measures and interventions with an impact on bacterial biofilms. *Agriculture*. 2022;12(8):1251. [DOI]
27. Castañeda-Gulla K, Sattlegger E, Mutukumira AN. Persistent contamination of *Salmonella*, *Campylobacter*, *Escherichia coli*, and *Staphylococcus aureus* at a broiler farm in New Zealand. *Canadian Journal of Microbiology*. 2020;66(3):171-85. [PMID: 31721603] [DOI]
28. Davies R, Wales A. Antimicrobial resistance on farms: a review including biosecurity and the potential role of disinfectants in resistance selection. *Comprehensive reviews in food science and food safety*. 2019;18(3):753-74. [PMID: 33336931] [DOI]
29. Salgueiro V, Manageiro V, Bandarra NM, Ferreira E, Clemente L, Caniça M. Genetic relatedness and diversity of *Staphylococcus aureus* from different reservoirs: humans and animals of livestock, poultry, zoo, and aquaculture. *Microorganisms*. 2020;8(9):1345. [PMID: 32899267] [PMCID: PMC7564200] [DOI]
30. Kozajda A, Ježak K, Kapsa A. Airborne *Staphylococcus aureus* in different environments—a review. *Environmental Science and Pollution Research*. 2019;26:34741-53. [PMID: 31654301] [PMCID: PMC6900272] [DOI]
31. Omidi AH, Sabati H, Amini S, Zonobian MA, Mohammadi MR. *Staphylococcus aureus* in the environment of healthcare centers. *Cell Mol Biomed Rep*. 2021;1(4):147-57. [DOI]
32. Wittekamp BH, Oostdijk EA, Cuthbertson BH, Brun-Buisson C, Bonten MJ. Selective decontamination of the digestive tract (SDD) in critically ill patients: a narrative review. *Intensive Care Medicine*. 2020;46(2):343-9. [PMID: 31820032] [PMCID: PMC7042187] [DOI]
33. Bi S, Westerholm M, Qiao W, Xiong L, Mahdy A, Yin D, et al. Metabolic performance of anaerobic digestion of chicken manure under wet, high solid, and dry conditions. *Bioresource technology*. 2020;296:122342. [PMID: 31711908] [DOI]
34. Boulianne M, Uzal FA, Opengart K. Clostridial diseases. *Diseases of Poultry*. 2020:966-94. [DOI]
35. Guo P, Zhang K, Ma X, He P. Clostridium species as probiotics: potentials and challenges. *Journal of animal science and biotechnology*. 2020;11:1-10. [PMID: 32099648] [PMCID: PMC7031906] [DOI]
36. Lorine D, Céline D, Frédéric B, Lorette H, Julie B, Laure M, et al. Influence of operating conditions on the persistence of *E. coli*, enterococci, *Clostridium perfringens* and *Clostridioides difficile* in semi-continuous mesophilic anaerobic reactors. *Waste Management*. 2021;134:32-41. [PMID: 34403994] [DOI]
37. Bindari YR, Kheravii SK, Morton CL, Wu S-B, Walkden-Brown SW, Gerber PF. Molecular detection of *Eimeria* species and *Clostridium perfringens* in poultry dust and pooled excreta of commercial broiler chicken flocks differing in productive performance. *Veterinary Parasitology*. 2021;291:109361. [PMID: 33550163] [DOI]
38. Fancher CA, Zhang L, Kiess AS, Adhikari PA, Dinh TT, Sukumaran AT. Avian pathogenic *Escherichia coli* and *Clostridium perfringens*: Challenges in no antibiotics ever broiler production and potential solutions. *Microorganisms*. 2020;8(10):1533. [PMID: 33036173] [PMCID: PMC7599686] [DOI]
39. Anju K, Karthik K, Divya V, Priyadarshini MLM, Sharma RK, Manoharan S. Toxinotyping and molecular characterization of antimicrobial resistance in *Clostridium perfringens* isolated from different sources of livestock and poultry. *Anaerobe*. 2021;67:102298. [PMID: 33220406] [DOI]
40. Abdolmohammadi Khiav L, Zahmatkesh A. Vaccination against pathogenic clostridia in animals: A review. *Tropical animal health and production*. 2021;53(2):284. [PMID: 33891221] [PMCID: PMC8062623] [DOI]
41. Gohari IM, Prescott JF. Clostridium. *Veterinary Microbiology*. 2022:309-34. [DOI]
42. Hustá M, Ducatelle R, Haesebrouck F, Van Immerseel F, Goossens E. A comparative study on the use of selective media for the enumeration of *Clostridium perfringens* in poultry faeces. *Anaerobe*. 2020;63:102205. [PMID: 32353484] [DOI]
43. Lepp D, Zhou Y, Ojha S, Mehdizadeh Gohari I, Carere J, Yang C, et al. Clostridium perfringens produces an adhesive pilus required for the pathogenesis of necrotic enteritis in poultry. *Journal of Bacteriology*. 2021;203(7):10.1128/jb.00578-20. [DOI]
44. Hussein EO, Ahmed SH, Abudabos AM, Suliman GM, Abd El-Hack ME, Swelum AA, et al. Ameliorative effects of antibiotic-, probiotic- and phytobiotic-supplemented diets on the performance, intestinal health, carcass traits, and meat quality of *Clostridium perfringens*-infected broilers. *Animals*. 2020;10(4):669. [PMID: 32290578] [PMCID: PMC7222811] [DOI]
45. Granstad S, Kristoffersen AB, Benestad SL, Sjurseth SK, David B, Sørensen L, et al. Effect of feed additives as alternatives to in-feed antimicrobials on production performance and intestinal *Clostridium perfringens* counts in broiler chickens. *Animals*. 2020;10(2):240. [PMID: 32028636] [PMCID: PMC7070674] [DOI]
46. Obianwuna UE, Qiu K, Wang J, Zhang H-j, Qi G-h, Huang L-l, et al. Effects of dietary Clostridium butyricum and fructooligosaccharides, alone or in combination, on performance, egg quality, amino acid digestibility, jejunal morphology, immune function, and antioxidant capacity of laying hens. *Frontiers in Microbiology*. 2023;14:1125897. [PMID: 36910205] [DOI]
47. Wang Y, Wang Y, Lin X, Gou Z, Fan Q, Jiang S. Effects of *Clostridium butyricum*, sodium butyrate, and butyric acid glycerides on the reproductive performance, egg quality, intestinal health, and offspring performance of yellow-feathered breeder hens. *Frontiers in*

- Microbiology. 2021;12:657542. [PMID: 34603221] [PMCID: PMC8481923] [DOI]
48. Niemi JK. The economic cost of bacterial infections. *Advancements and technologies in pig and poultry bacterial disease control*. 2021;1-23. [DOI]
49. Igbinosa EO, Beshiru A, Igbinosa IH, Ogofure AG, Ekundayo TC, Okoh AI. Prevalence, multiple antibiotic resistance and virulence profile of methicillin-resistant *Staphylococcus aureus* (MRSA) in retail poultry meat from Edo, Nigeria. *Frontiers in Cellular and Infection Microbiology*. 2023;13:1122059. [PMID: 36936767] [PMCID: PMC10017849] [DOI]
50. Ogundipe FO, Ojo OE, Feßler AT, Hanke D, Awoyomi OJ, Ojo DA, et al. Antimicrobial resistance and virulence of methicillin-resistant *Staphylococcus aureus* from human, chicken and environmental samples within live bird markets in three Nigerian cities. *Antibiotics*. 2020;9(9):588. [PMID: 32911712] [PMCID: PMC7558163] [DOI]
51. Benrabia I, Hamdi TM, Shehata AA, Neubauer H, Wareth G. Methicillin-resistant *Staphylococcus aureus* (MRSA) in poultry species in Algeria: Long-term study on prevalence and antimicrobial resistance. *Veterinary sciences*. 2020;7(2):54. [PMID: 32349228] [PMCID: PMC7356745] [DOI]
52. Okorie-Kanu OJ, Anyanwu MU, Ezenduka EV, Mgbeahuruike AC, Thapaliya D, Gerbig G, et al. Molecular epidemiology, genetic diversity and antimicrobial resistance of *Staphylococcus aureus* isolated from chicken and pig carcasses, and carcass handlers. *PLoS One*. 2020;15(5):e0232913. [PMID: 32407414] [PMCID: PMC7224487] [DOI]
53. Saharan VV, Verma P, Singh AP. *Escherichia coli*, *Salmonella* spp., and *Staphylococcus aureus* susceptibility to antimicrobials of human and veterinary importance in poultry sector of India. *Journal of food safety*. 2020;40(1):e12742. [DOI]
54. Gržinić G, Piotrowicz-Cieślak A, Klimkiewicz-Pawlas A, Gómy RL, Ławniczek-Walczak A, Piechowicz L, et al. Intensive poultry farming: A review of the impact on the environment and human health. *Science of the Total Environment*. 2023;858:160014. [PMID: 36368402] [DOI]
55. Assafi MS, Hado HA, Abdulrahman IS. Detection of methicillin-resistant *Staphylococcus aureus* in broiler and broilers farm workers in Duhok, Iraq by using conventional and PCR techniques. *Iraqi J Vet Sci*. 2020;34(1):15-22. [DOI]
56. Marcon AV, De Oliveira G, Caldara FR, Garcia RG, Matins R, Marcon A, et al. Bacteriological and histopathological evaluation of articulations of chickens diagnosed with arthritis. *Brazilian Journal of Poultry Science*. 2019;21. [DOI]
57. Ekesi NS. Examining Pathogenesis and Preventatives in Spontaneous and *Staphylococcus*-Induced Bacterial Chondronecrosis with Osteomyelitis in Broilers: University of Arkansas; 2020.
58. Tabar MM, Peighambari SM, Bagheri S. Lameness caused by *Staphylococcus aureus* in poultry: A review. *The Journal of Poultry Sciences and Avian Diseases*. 2024;2(1):11-7. [DOI]
59. Logue CM, Andreasen CB, Borst LB, Eriksson H, Hampson DJ, Sanchez S, et al. Other bacterial diseases. *Diseases of poultry*. 2020:995-1085. [DOI]
60. Chen C, Wu F. Livestock-associated methicillin-resistant *Staphylococcus aureus* (LA-MRSA) colonisation and infection among livestock workers and veterinarians: a systematic review and meta-analysis. *Occupational and Environmental Medicine*. 2021;78(7):530-40. [PMID: 33097674] [DOI]
61. Crespo-Piazuelo D, Lawlor PG. Livestock-associated methicillin-resistant *Staphylococcus aureus* (LA-MRSA) prevalence in humans in close contact with animals and measures to reduce on-farm colonisation. *Irish veterinary journal*. 2021;74:1-12. [PMID: 34362463] [PMCID: PMC8348836] [DOI]
62. Fetsch A, Etter D, Johler S. Livestock-associated methicillin-resistant *Staphylococcus aureus*—Current situation and impact from a One Health perspective. *Current Clinical Microbiology Reports*. 2021;8(3):103-13. [DOI]
63. Liene T, Schnitt A, Cuny C, Maurischat S, Tenhagen B-A. Phylogenetic tracking of LA-MRSA ST398 intra-farm transmission among animals, humans and the environment on German dairy farms. *Microorganisms*. 2021;9(6):1119. [PMID: 34064246] [PMCID: PMC8224388] [DOI]
64. Silva V, Araújo S, Monteiro A, Eira J, Pereira JE, Maltez L, et al. *Staphylococcus aureus* and MRSA in livestock: antimicrobial resistance and genetic lineages. *Microorganisms*. 2023;11(1):124. [PMID: 36677414] [PMCID: PMC9865216] [DOI]
65. Silva V, Vieira-Pinto M, Saraiva C, Manageiro V, Reis L, Ferreira E, et al. Prevalence and characteristics of multidrug-resistant livestock-associated methicillin-resistant *Staphylococcus aureus* (LA-MRSA) CC398 isolated from quails (*Coturnix Coturnix Japonica*) slaughtered for human consumption. *Animals*. 2021;11(7):2038. [PMID: 34359166] [PMCID: PMC8300319] [DOI]
66. Dombrowski S. *Berichte zur Lebensmittelsicherheit 2012: Monitoring*: Springer-Verlag; 2013.
67. Seker E, Ozenc E, Turedi OK, Yilmaz M. Prevalence of *mecA* and *pvl* genes in coagulase negative staphylococci isolated from bovine mastitis in smallholder dairy farms in Turkey. *Animal Biotechnology*. 2023;34(7):2427-32. [PMID: 35792781] [DOI]
68. Franceschini G, Bottino M, Millet I, Martello E, Zaltron F, Favretto AR, et al. Assessment of the exposure of turkey farmers to antimicrobial resistance associated with working practices. *Veterinary sciences*. 2019;6(1):13. [PMID: 30717204] [PMCID: PMC6466403] [DOI]
69. Hafez H, Shehata A. Turkey production and health: Current challenges. *Ger J Vet Res*. 2021;1(1):3-14. [DOI]
70. Saadati A, Mashak Z, Yarmand MS. Prevalence of staphylococcal cassette chromosome *mec* and *panton-valentine* leukocidin gene amongst the methicillin-resistant *Staphylococcus aureus* strains isolated from fowl meat. *Int J Enteric Pathog*. 2019;7:93-8. [DOI]
71. für Risikobewertung B. *Erreger von Zoonosen in Deutschland im Jahr...: 2014*: BfR, Pressestelle; 2016.
72. Sheet OH, Hussein SA, Al-Chalaby AY. Detection of methicillin-resistant *Staphylococcus aureus* from broiler carcasses in Mosul city. *Iraqi Journal of Veterinary Sciences*. 2021;35(3):489-93. [DOI]
73. Belhout C, Elgroud R, Butaye P. Methicillin-Resistant *Staphylococcus aureus* (MRSA) and other methicillin-resistant staphylococci and *Mammaliococcus* (MRNaS) associated with animals and food products in Arab countries: A review. *Veterinary Sciences*. 2022;9(7):317. [PMID: 35878334] [PMCID: PMC9320237] [DOI]
74. Elsayed MM, Elkenany RM, El-Khateeb AY, Nabil NM, Tawakol MM, Hassan HM. Isolation and encapsulation of bacteriophage with chitosan nanoparticles for biocontrol of multidrug-resistant methicillin-resistant *Staphylococcus aureus* isolated from broiler poultry farms. *Scientific Reports*. 2024;14(1):4702. [PMID: 38409454] [PMCID: PMC10897325] [DOI]
75. Ralte R, Kumar V, Wani MY, Randhawa S, Malik A, Kaur N. Antimicrobial resistance profile of livestock and poultry of North-western Punjab. *Int J Curr Microbiol Appl Sci*. 2019;8(10):1250-9. [DOI]
76. Ramandinianto SC, Khairullah AR, Effendi MH, Hestiana EP. Profile of multidrug resistance (MDR) and methicillin resistant *Staphylococcus aureus* (MRSA) on dairy farms in east Java province, Indonesia. *Indian Journal of Forensic Medicine & Toxicology*. 2020;14(4):3439-45. [DOI]
77. Hamza M, Sivaraman G, Mothadaka MP. Evolution, Characteristics, and Clonal Expansion of Livestock-Associated Methicillin-Resistant *Staphylococcus aureus* (LA-MRSA): Global Perspectives. *Handbook on Antimicrobial Resistance: Current Status,*

- Trends in Detection and Mitigation Measures: Springer; 2023. p. 1-20[DOI]
78. Parvin MS, Ali MY, Talukder S, Nahar A, Chowdhury EH, Rahman MT, et al. Prevalence and multidrug resistance pattern of methicillin resistant *S. aureus* isolated from frozen chicken meat in Bangladesh. *Microorganisms*. 2021;9(3):636. [PMID: 33803779] [PMCID: PMC8003192] [DOI]
 79. Khodary M, Nasr El-Deen N, Gamal El-Deen I. Effect of experimental *Clostridium perfringens* infection on some immunological, hematological and biochemical values in broiler chickens. *Zagazig Veterinary Journal*. 2019;47(2):222-33. [DOI]
 80. Mora ZV-dl, Macías-Rodríguez ME, Arratia-Quijada J, Gonzalez-Torres YS, Nuño K, Villarruel-López A. *Clostridium perfringens* as foodborne pathogen in broiler production: pathophysiology and potential strategies for controlling necrotic enteritis. *Animals*. 2020;10(9):1718. [PMID: 32972009] [PMCID: PMC7552638] [DOI]
 81. Nicholds J, McQuain C, Hofacre C, Mathis G, Fuller A, Telg B, et al. The effect of different species of *Eimeria* with *Clostridium perfringens* on performance parameters and induction of clinical necrotic enteritis in broiler chickens. *Avian diseases*. 2021;65(1):132-7. [PMID: 34339132] [DOI]
 82. Abd El-Hack ME, El-Saadony MT, Elbestawy AR, Nahed A, Saad AM, Salem HM, et al. Necrotic enteritis in broiler chickens: disease characteristics and prevention using organic antibiotic alternatives—a comprehensive review. *Poultry science*. 2022;101(2):101590. [PMID: 34953377] [PMCID: PMC8715378] [DOI]
 83. Emami NK, Dalloul RA. Centennial Review: Recent developments in host-pathogen interactions during necrotic enteritis in poultry. *Poultry Science*. 2021;100(9):101330. [PMID: 34280643] [PMCID: PMC8318987] [DOI]
 84. Fathima S, Hakeem WGA, Shanmugasundaram R, Selvaraj RK. Necrotic enteritis in broiler chickens: A review on the pathogen, pathogenesis, and prevention. *Microorganisms*. 2022;10(10):1958. [PMID: 36296234] [PMCID: PMC9610872] [DOI]
 85. Fancher C. Prevalence and Molecular Characteristics of Avian Pathogenic *Escherichia coli* and *Clostridium perfringens* in 'No Antibiotics Ever' Broiler Farms: Mississippi State University; 2021.
 86. Shelke P, Pawade M, Mhase P, Mote C, Lonkar V, Suryawanshi D, et al. Toxinotyping of *Clostridium perfringens* in Poultry from Necrotic Enteritis Cases. *Journal of Animal Research*. 2020;10(1):149-55. [DOI]
 87. Helal SS, Khalaf NM, El Menisy AA, Lebdah MA. *Clostridium perfringens* type A causing necrotic enteritis outbreaks among chickens in Egypt. *Zagazig Veterinary Journal*. 2019;47(4):398-407. [DOI]
 88. Praveen Kumar N, Vinod Kumar N, Karthik A. Molecular detection and characterization of *Clostridium perfringens* toxin genes causing necrotic enteritis in broiler chickens. *Tropical animal health and production*. 2019;51(6):1559-69. [PMID: 31076994] [DOI]
 89. Grenda T, Jarosz A, Sapała M, Grenda A, Patyra E, Kwiatek K. *Clostridium perfringens*—opportunistic Foodborne Pathogen, its diversity and epidemiological significance. *Pathogens*. 2023;12(6):768. [PMID: 37375458] [PMCID: PMC10304509] [DOI]
 90. Calik A, Omara II, White MB, Evans NP, Kamezos TP, Dalloul RA. Dietary non-drug feed additive as an alternative for antibiotic growth promoters for broilers during a necrotic enteritis challenge. *Microorganisms*. 2019;7(8):257. [PMID: 31412542] [PMCID: PMC6723652] [DOI]
 91. Gharib-Naseri K, Kheravii S, Keerqin C, Morgan N, Swick R, Choct M, et al. Two different *Clostridium perfringens* strains produce different levels of necrotic enteritis in broiler chickens. *Poultry science*. 2019;98(12):6422-32. [PMID: 31424518] [PMCID: PMC8913766] [DOI]
 92. Khalique A, Zeng D, Shoaib M, Wang H, Qing X, Rajput DS, et al. Probiotics mitigating subclinical necrotic enteritis (SNE) as potential alternatives to antibiotics in poultry. *Amb Express*. 2020;10:1-10. [PMID: 32172398] [PMCID: PMC7072080] [DOI]
 93. Mwangi S, Timmons J, Fitz-Coy S, Parveen S. Characterization of *Clostridium perfringens* recovered from broiler chicken affected by necrotic enteritis. *Poultry Science*. 2019;98(1):128-35. [PMID: 30053181] [DOI]
 94. Calvert AJ. The effects of *Bacillus subtilis*, dietary calcium and coccidial vaccines in broilers experiencing naturally occurring necrotic enteritis. 2022.
 95. Cardoso Dal Pont G, Farnell M, Farnell Y, Kogut MH. Dietary factors as triggers of low-grade chronic intestinal inflammation in poultry. *Microorganisms*. 2020;8(1):139. [PMID: 31963876] [PMCID: PMC7022292] [DOI]
 96. Kulkarni RR, Gaghan C, Gorrell K, Sharif S, Taha-Abdelaziz K. Probiotics as alternatives to antibiotics for the prevention and control of necrotic enteritis in chickens. *Pathogens*. 2022;11(6):692. [PMID: 35745546] [PMCID: PMC9229159] [DOI]
 97. Rajput DS, Zeng D, Khalique A, Rajput SS, Wang H, Zhao Y, et al. Pretreatment with probiotics ameliorate gut health and necrotic enteritis in broiler chickens, a substitute to antibiotics. *Amb Express*. 2020;10:1-11. [PMID: 33336284] [PMCID: PMC7746796] [DOI]
 98. Dong Q, Liu Y, Li W, Liu Y, Ye X. Cross-species transmission risk of livestock-associated MRSA: A systematic review and Bayesian meta-analysis of global data. *Preventive Veterinary Medicine*. 2021;194:105429. [PMID: 34271475] [DOI]
 99. da Silva AC, Rodrigues MX, Silva NCC. Methicillin-resistant *Staphylococcus aureus* in food and the prevalence in Brazil: a review. *Brazilian Journal of Microbiology*. 2020;51(1):347-56. [PMID: 31667799] [PMCID: PMC7058716] [DOI]
 100. Barcudi D, Sosa EJ, Lamberghini R, Garnerio A, Tosoroni D, Decca L, et al. MRSA dynamic circulation between the community and the hospital setting: New insights from a cohort study. *Journal of Infection*. 2020;80(1):24-37. [PMID: 31606351] [DOI]
 101. Hossain MJ, Sohidullah M, Alam MA, Al Mamun MS, Badr Y, Altaib H, et al. Molecular Detection of Methicillin Resistant *Staphylococcus aureus* (MRSA) in Poultry in Bangladesh: Having Public Health Significance. *European Journal of Veterinary Medicine*. 2022;2(6):17-21. [DOI]
 102. Andrew Seleledi L, Mohammed Hassan Z, Manyelo TG, Mabelebele M. The current status of the alternative use to antibiotics in poultry production: An African perspective. *Antibiotics*. 2020;9(9):594. [PMID: 32932751] [PMCID: PMC7559726] [DOI]
 103. Samad A. Antibiotics resistance in poultry and its solution. *Devotion: Journal of Research and Community Service*. 2022;3(10):999-1020. [DOI]
 104. Amoako DG, Somboro AM, Abia AL, Molechan C, Perrett K, Bester LA, et al. Antibiotic resistance in *Staphylococcus aureus* from poultry and poultry products in uMgungundlovu District, South Africa, using the "Farm to Fork" approach. *Microbial Drug Resistance*. 2020;26(4):402-11. [PMID: 31647362] [DOI]
 105. Kwoji ID, Jauro S, Musa JA, Lekko YM, Salihu SI, Danchuwa HA. Phenotypic detection of methicillin-resistant *Staphylococcus aureus* in village chickens from poultry markets in Maiduguri, Nigeria. *Journal of Advanced Veterinary and Animal Research*. 2019;6(2):163. [PMID: 31453186] [PMCID: PMC6702879] [DOI]
 106. Peter I, Ngwu J, Edemekong C, Ugwueke I, Uzoeto H, Joseph O, et al. First report prevalence of livestock acquired Methicillin resistant *Staphylococcus aureus* (LA-MRSA) strain in south eastern, Nigeria. *IOSR J Nurs Health Sci*. 2022;11(1):50-6.
 107. Craft KM, Nguyen JM, Berg LJ, Townsend SD. Methicillin-resistant *Staphylococcus aureus* (MRSA): antibiotic-resistance and the biofilm phenotype. *MedChemComm*. 2019;10(8):1231-41. [PMID: 31534648] [PMCID: PMC6748282] [DOI]

108. Persoons D, Van Hoorebeke S, Hermans K, Butaye P, De Kruijff A, Haesebrouck F, et al. Methicillin-resistant *Staphylococcus aureus* in poultry. *Emerging infectious diseases*. 2009;15(3):452. [PMID: 19239762] [PMCID: PMC2666289] [DOI]
109. Shareef A, Mansour R, Ibrahim K. *Staphylococcus aureus* in commercial breeder layer flocks. *Iraqi Journal of Veterinary Sciences*. 2009;23(3).
110. Saud B, Paudel G, Khichaju S, Bajracharya D, Dhungana G, Awasthi MS, et al. Multidrug-resistant bacteria from raw meat of buffalo and chicken, Nepal. *Veterinary medicine international*. 2019;2019(1):7960268. [PMID: 31186828] [PMCID: PMC6521380] [DOI]
111. Zehra A, Gulzar M, Singh R, Kaur S, Gill J. Prevalence, multidrug resistance and molecular typing of methicillin-resistant *Staphylococcus aureus* (MRSA) in retail meat from Punjab, India. *Journal of global antimicrobial resistance*. 2019;16:152-8. [PMID: 30312831] [DOI]
112. Lin Q, Sun H, Yao K, Cai J, Ren Y, Chi Y. The prevalence, antibiotic resistance and biofilm formation of *Staphylococcus aureus* in bulk ready-to-eat foods. *Biomolecules*. 2019;9(10):524. [PMID: 31548510] [PMCID: PMC6843627] [DOI]
113. Mekhloufi OA, Chieffi D, Hammoudi A, Bensefia SA, Fanelli F, Fusco V. Prevalence, enterotoxigenic potential and antimicrobial resistance of *Staphylococcus aureus* and methicillin-resistant *Staphylococcus aureus* (MRSA) isolated from Algerian ready to eat foods. *Toxins*. 2021;13(12):835. [PMID: 34941673] [PMCID: PMC8707561] [DOI]
114. Mesbah A, Mashak Z, Abdolmaleki Z. A survey of prevalence and phenotypic and genotypic assessment of antibiotic resistance in *Staphylococcus aureus* bacteria isolated from ready-to-eat food samples collected from Tehran Province, Iran. *Tropical Medicine and Health*. 2021;49:1-12. [PMID: 34635183] [PMCID: PMC8504009] [DOI]
115. Lika E, Puvača N, Jeremić D, Stanojević S, Shtylla Kika T, Cocoli S, et al. Antibiotic susceptibility of *Staphylococcus* species isolated in raw chicken meat from retail stores. *Antibiotics*. 2021;10(8):904. [PMID: 34438954] [PMCID: PMC8388630] [DOI]
116. Jackson KA, Gokhale RH, Nadle J, Ray SM, Dumyati G, Schaffner W, et al. Public health importance of invasive methicillin-sensitive *Staphylococcus aureus* infections: surveillance in 8 US counties, 2016. *Clinical Infectious Diseases*. 2020;70(6):1021-8. [PMID: 31245810] [PMCID: PMC7902232] [DOI]
117. Miranda C, Silva V, Igrejas G, Poeta P. Impact of European pet antibiotic use on enterococci and staphylococci antimicrobial resistance and human health. *Future Microbiology*. 2021;16(3):185-203. [PMID: 33491483] [DOI]
118. Park S, Ronholm J. *Staphylococcus aureus* in agriculture: lessons in evolution from a multispecies pathogen. *Clinical microbiology reviews*. 2021;34(2):10.1128/cmr.00182-20. [PMID: 33568553] [PMCID: PMC7950364] [DOI]
119. Samutela MT, Kwenda G, Simulundu E, Nkhoma P, Higashi H, Frey A, et al. Pigs as a potential source of emerging livestock-associated *Staphylococcus aureus* in Africa: a systematic review. *International Journal of Infectious Diseases*. 2021;109:38-49. [PMID: 34146692] [DOI]
120. Tao C-W, Chen J-S, Hsu B-M, Koner S, Hung T-C, Wu H-M, et al. Molecular evaluation of traditional chicken farm-associated bioaerosols for methicillin-resistant *Staphylococcus aureus* shedding. *Antibiotics*. 2021;10(8):917. [PMID: 34438967] [PMCID: PMC8388662] [DOI]
121. Kozajda A, Miśkiewicz E, Jeżak K. Zoonotic bacteria in the vicinity of animal farms as a factor disturbing the human microbiome: a review. *International Journal of Occupational Medicine and Environmental Health*. 2024;37(2):138. [PMID: 38577723] [PMCID: PMC1142397] [DOI]
122. García-Vela S, Martínez-Sancho A, Said LB, Torres C, Fliss I. Pathogenicity and Antibiotic Resistance Diversity in *Clostridium perfringens* Isolates from Poultry Affected by Necrotic Enteritis in Canada. *Pathogens*. 2023;12(7):905. [PMID: 37513752] [PMCID: PMC10383762] [DOI]
123. Wei B, Cha S-Y, Zhang J-F, Shang K, Park H-C, Kang J, et al. Antimicrobial susceptibility and association with toxin determinants in *Clostridium perfringens* isolates from chickens. *Microorganisms*. 2020;8(11):1825. [PMID: 33228100] [PMCID: PMC7699427] [DOI]
124. Aljumaah MR, Alkhulaifi MM, Abudabos AM. In vitro antibacterial efficacy of non-antibiotic growth promoters in poultry industry. *The Journal of Poultry Science*. 2020;57(1):45-54. [PMID: 32174764] [PMCID: PMC7063082] [DOI]
125. Hussein EO, Ahmed SH, Abudabos AM, Aljumaah MR, Alkhulaifi MM, Nassan MA, et al. Effect of antibiotic, phytobiotic and probiotic supplementation on growth, blood indices and intestine health in broiler chicks challenged with *Clostridium perfringens*. *Animals*. 2020;10(3):507. [PMID: 32197455] [PMCID: PMC7142990] [DOI]
126. Rafiq K, Tofazzal Hossain M, Ahmed R, Hasan MM, Islam R, Hossen MI, et al. Role of different growth enhancers as alternative to in-feed antibiotics in poultry industry. *Frontiers in veterinary science*. 2022;8:794588. [PMID: 35224074] [PMCID: PMC8873819] [DOI]
127. Agunos A, Léger DF, Carson CA, Gow SP, Bosman A, Irwin RJ, et al. Antimicrobial use surveillance in broiler chicken flocks in Canada, 2013-2015. *PloS one*. 2017;12(6):e0179384. [PMID: 28658278] [PMCID: PMC5489168] [DOI]
128. MacDonald JM, Wang SL. Foregoing sub-therapeutic antibiotics: the impact on broiler grow-out operations. *Applied Economic Perspectives and Policy*. 2011;33(1):79-98. [DOI]
129. Teillant A, Laxminarayan R. Economics of antibiotic use in US swine and poultry production. *Choices*. 2015;30(1):1-11.
130. Iannetti L, Romagnoli S, Cotturone G, Podaliri Vulpiani M. Animal welfare assessment in antibiotic-free and conventional broiler chicken. *Animals*. 2021;11(10):2822. [PMID: 34679843] [PMCID: PMC8532607] [DOI]
131. Singer RS, Porter LJ, Schrag NF, Davies PR, Apley MD, Bjork K. Estimates of on-farm antimicrobial usage in broiler chicken production in the United States, 2013–2017. *Zoonoses and Public Health*. 2020;67:22-35. [DOI]
132. Singer RS, Porter LJ, Thomson DU, Gage M, Beaudoin A, Wishnie JK. Raising animals without antibiotics: US producer and veterinarian experiences and opinions. *Frontiers in veterinary science*. 2019;6:452. [PMID: 31867349] [PMCID: PMC6910073] [DOI]
133. AlJindan R, AlEraky DM, Farhat M, Almandil NB, AbdulAzeez S, Borgio JF. Genomic insights into virulence factors and multi-drug resistance in *Clostridium perfringens* IRMC2505A. *Toxins*. 2023;15(6):359. [PMID: 37368661] [PMCID: PMC10302382] [DOI]
134. Bendary MM, Abd El-Hamid MI, El-Tarabili RM, Hefny AA, Algendy RM, Elzohairy NA, et al. *Clostridium perfringens* associated with foodborne infections of animal origins: Insights into prevalence, antimicrobial resistance, toxin genes profiles, and toxinotypes. *Biology*. 2022;11(4):551. [PMID: 35453750] [PMCID: PMC9028928] [DOI]
135. Li J, Zhou Y, Yang D, Zhang S, Sun Z, Wang Y, et al. Prevalence and antimicrobial susceptibility of *Clostridium perfringens* in chickens and pigs from Beijing and Shanxi, China. *Veterinary Microbiology*. 2021;252:108932. [PMID: 33316633] [DOI]
136. Ali M, Islam M. Characterization of β -lactamase and quinolone resistant *Clostridium perfringens* recovered from broiler chickens with necrotic enteritis in Bangladesh. *Iranian journal of veterinary research*. 2021;22(1):48.
137. Mohiuddin M, Song Z, Liao S, Qi N, Li J, Lv M, et al. Animal model studies, antibiotic resistance and toxin gene profile of ne-reproducing *Clostridium perfringens* type A and type G strains isolated from commercial poultry farms in China. *Microorganisms*. 2023;11(3):622. [PMID: 36985195] [PMCID: PMC10059142] [DOI]

138. Xu W, Wang H, Liu L, Miao Z, Huo Y, Zhong Z. Prevalence and characterization of *Clostridium perfringens* isolated from different chicken farms in China. *Anaerobe*. 2021;72:102467. [PMID: 34688908] [DOI]
139. Hedman HD, Vasco KA, Zhang L. A review of antimicrobial resistance in poultry farming within low-resource settings. *Animals*. 2020;10(8):1264. [PMID: 32722312] [PMCID: PMC7460429] [DOI]
140. Kumar D, Pornsukarom S, Thakur S. Antibiotic usage in poultry production and antimicrobial-resistant *Salmonella* in poultry. *Food safety in poultry meat production*. 2019:47-66. [DOI]
141. Jang Y-S, Kim D-H, Bae D, Kim S-H, Kim H, Moon J-S, et al. Prevalence, toxin-typing, and antimicrobial susceptibility of *Clostridium perfringens* from retail meats in Seoul, Korea. *Anaerobe*. 2020;64:102235. [PMID: 32619505] [DOI]
142. Bortoluzzi C, Serpa Vieira B, de Paula Dorigam JC, Menconi A, Sokale A, Doranalli K, et al. *Bacillus subtilis* DSM 32315 supplementation attenuates the effects of *Clostridium perfringens* challenge on the growth performance and intestinal microbiota of broiler chickens. *Microorganisms*. 2019;7(3):71. [PMID: 30841654] [PMCID: PMC6463140] [DOI]
143. Sandvang D, Skjoet-Rasmussen L, Cantor MD, Mathis GF, Lumpkins BS, Blanch A. Effects of feed supplementation with 3 different probiotic *Bacillus* strains and their combination on the performance of broiler chickens challenged with *Clostridium perfringens*. *Poultry Science*. 2021;100(4):100982. [PMID: 33647715] [PMCID: PMC7921869] [DOI]
144. Sun Y, Ni A, Jiang Y, Li Y, Huang Z, Shi L, et al. Effects of replacing in-feed antibiotics with synergistic organic acids on growth performance, health, carcass, and immune and oxidative statuses of broiler chickens under *clostridium perfringens* type a challenge. *Avian Diseases*. 2020;64(3):393-400. [PMID: 33205169] [DOI]
145. Brusseau M, Ramirez-Andreotta M, Pepper I, Maximillian J. Environmental impacts on human health and well-being. *Environmental and pollution science*: Elsevier; 2019. p. 477-99 [DOI]
146. Mkangara M. Prevention and control of human *Salmonella enterica* infections: An implication in food safety. *International Journal of Food Science*. 2023;2023(1):8899596. [PMID: 37727836] [PMCID: PMC10506869] [DOI]
147. Vajda Á, Ózsvári L, Szakos D, Kasza G. Estimation of the impact of foodborne salmonellosis on consumer well-being in Hungary. *International Journal of Environmental Research and Public Health*. 2021;18(19):10131. [PMID: 34639433] [PMCID: PMC8507706] [DOI]
148. Gallo M, Ferrara L, Calogero A, Montesano D, Naviglio D. Relationships between food and diseases: What to know to ensure food safety. *Food Research International*. 2020;137:109414. [PMID: 33233102] [DOI]
149. Mahros MA, Abd-Elghany SM, Sallam KI. Multidrug-, methicillin-, and vancomycin-resistant *Staphylococcus aureus* isolated from ready-to-eat meat sandwiches: An ongoing food and public health concern. *International Journal of Food Microbiology*. 2021;346:109165. [PMID: 33770679] [DOI]
150. Silva V, Caniça M, Ferreira E, Vieira-Pinto M, Saraiva C, Pereira JE, et al. Multidrug-resistant methicillin-resistant coagulase-negative staphylococci in healthy poultry slaughtered for human consumption. *Antibiotics*. 2022;11(3):365. [PMID: 35326828] [PMCID: PMC8944763] [DOI]
151. Cuny C, Layer F, Hansen S, Werner G, Witte W. Nasal colonization of humans with occupational exposure to raw meat and to raw meat products with methicillin-susceptible and methicillin-resistant *Staphylococcus aureus*. *Toxins*. 2019;11(4):190. [PMID: 30935022] [PMCID: PMC6521318] [DOI]
152. Thwala T, Madoroba E, Basson A, Butaye P. Prevalence and characteristics of *Staphylococcus aureus* associated with meat and meat products in African countries: a review. *Antibiotics*. 2021;10(9):1108. [PMID: 34572690] [PMCID: PMC8465003] [DOI]
153. Jangra S, Khunger S, Chattopadhyaya D. Antimicrobial Resistance in *Staphylococci* Special Emphasis on Methicillin Resistance among Companion Livestock and Its Impact on Human Health in Rural India. *Bovine Science-Challenges and Advances*: IntechOpen; 2022 [DOI]
154. González-Machado C, Alonso-Calleja C, Capita R. Prevalence and types of methicillin-resistant *Staphylococcus aureus* (MRSA) in meat and meat products from retail outlets and in samples of animal origin collected in farms, slaughterhouses and meat processing facilities. A review. *Food Microbiology*. 2024:104580. [PMID: 39038886] [DOI]
155. Ahmed R, Fathy M, Al-Azeem A, Mohamed W, Elghoneimy A, Mohamed I. *Staphylococcus aureus* in poultry: prevalence and antibiogram of methicillin-resistant *Staphylococcus aureus* in avian species in the Southern Provinces of Egypt. *Egyptian Journal of Microbiology*. 2021;56(1):89-98. [DOI]
156. Colelli G, Normanno G. Methicillin Resistant *Staphylococcus aureus* (MRSA) in raw buffalo milk and its fate along the human gastrointestinal tract: an in vitro study. 2020.
157. González-Machado C, Capita R, Alonso-Calleja C. Methicillin-Resistant *Staphylococcus aureus* (MRSA) in Dairy Products and Bulk-Tank Milk (BTM). *Antibiotics*. 2024;13(7):588. [PMID: 39061270] [PMCID: PMC11273636] [DOI]
158. Khairullah AR, Kurniawan SC, Effendi MH, Sudjarwo SA, Ramandinianto SC, Widodo A, et al. A review of new emerging livestock-associated methicillin-resistant *Staphylococcus aureus* from pig farms. *Veterinary World*. 2023;16(1):46. [PMID: 36855358] [PMCID: PMC9967705] [DOI]
159. Sneeringer S, MacDonald JM, Key N, McBride WD, Mathews K. Economics of antibiotic use in US livestock production. *USDA, Economic Research Report*. 2015(200).
160. Hamad GM, Abdelmotilib NM, Darwish AM, Zeitoun AM. Commercial probiotic cell-free supernatants for inhibition of *Clostridium perfringens* poultry meat infection in Egypt. *Anaerobe*. 2020;62:102181. [PMID: 32092413] [DOI]
161. Mohammadi TN, Shen C, Li Y, Zayda MG, Sato J, Masuda Y, et al. Characterization of *Clostridium perfringens* bacteriophages and their application in chicken meat and milk. *International Journal of Food Microbiology*. 2022;361:109446. [PMID: 34742146] [DOI]
162. García S, Vidal JE, Heredia N, Juneja VK. *Clostridium perfringens*. *Food Microbiology: Fundamentals and Frontiers*. 2019:513-40. [PMID: PMC7357358] [DOI]
163. Fu Y, Alenezi T, Sun X. *Clostridium perfringens*-induced necrotic diseases: an overview. *Immuno*. 2022;2(2):387-407. [DOI]
164. OLUSEYI DF. ISOLATION AND ANTIBIOTICS SUSCEPTIBILITY OF *CLOSTRIDIUM PERFRINGENS*. 2022.
165. Priya GB, Srinivas K, Shilla H, Milton AAP. High Prevalence of Multidrug-Resistant, Biofilm-Forming Virulent *Clostridium perfringens* in Broiler Chicken Retail Points in Northeast India. *Foods*. 2023;12(22):4185. [PMID: 38002242] [PMCID: PMC10670561] [DOI]
166. Zheng X, Teng D, Mao R, Hao Y, Yang N, Hu F, et al. A study on fungal defensin against multidrug-resistant *Clostridium perfringens* and its treatment on infected poultry. *Applied Microbiology and Biotechnology*. 2021;105:7265-82. [PMID: 34491399] [DOI]